

## Uniform Approximation of Differentiable Functions by Algebraic Polynomials

HERBERT F. SINWEL\*

*Mathematisches Institut, Johannes Kepler Universität,  
Linz, Austria*

*Communicated by P. L. Butzer*

Received June 7, 1979

### INTRODUCTION

In this paper we deduce a special form of Jackson's fundamental direct theorem of best approximation. We give an asymptotic best upper bound of the uniform approximation error of differentiable functions by algebraic polynomials.

We attempt to find the algebraic version of the theorem of Favard [1], and Achieser and Krein [2]:

$$E_n(W_r) = K_r(n+1)^{-r}. \quad (1)$$

In Sections 2 and 3 we prove the following

**THEOREM 1.** *Let  $r \in \mathbb{N}$  and  $n \geq r - 1$ . Then*

$$E_n(W_r) < K_r \cdot (n+1-r)! / (n+1)! \quad (2)$$

The constant  $K_r$  cannot be improved. This fact is a result of Bernstein's theorem [4]:

$$\lim_{n \rightarrow \infty} n^r \cdot E_n(W_r) = K_r. \quad (3)$$

The proofs of (1) and (3) can be found in [5].

\* Research supported by the Austrian Fonds zur Förderung der wissenschaftlichen Forschung.

Bernstein's theorem also implies the existence of constants  $L_r$  such that for all  $r \in \mathbb{N}$  and  $n \geq r - 1$ ,

$$\int E_n(W_r) \leq L_r \cdot n^{-r}; \quad (4)$$

but it does not give upper bounds for  $L_r$ .

## 1. DEFINITIONS

Let  $C[-1, 1]$  be the space of all real continuous functions on  $[-1, 1]$  with the norm

$$\|g\| := \max_{-1 \leq x \leq 1} |g(x)|.$$

Let  $\mathcal{P}_n$  be the space of all algebraic polynomials of degree at most  $n$ ; furthermore, let

$$E_n(g) := \inf_{p \in \mathcal{P}_n} \|g - p\|$$

be the approximation error on  $C[-1, 1]$ . Let  $W_r$  be the space of all functions  $g$  with  $g, g', g'', \dots, g^{(r-1)}$  in  $C[-1, 1]$  and  $|g^{(r)}| \leq 1$  a.e. The constant  $E_n(W_r)$  is defined by

$$E_n(W_r) := \sup_{g \in W_r} E_n(g).$$

Analogously let  $C_{2\pi}$  be the space of all real continuous  $2\pi$ -periodic functions with the norm ( $K := [\theta, 2\pi]$ )

$$\|f\|^* := \max_{x \in K} |f(x)|.$$

Let  $\mathcal{E}_n$  be the space of all trigonometric polynomials of degree at most  $n$  and let

$$E_n^*(f) := \inf_{t \in \mathcal{E}_n} \|f - t\|^*$$

be the approximation error on  $C_{2\pi}$ . Let  $W_r^*$  be the space of all functions  $f$  with  $f, f', f'', \dots, f^{(r-1)}$  in  $C_{2\pi}$  and  $|f^{(r)}| \leq 1$  a.e. The constant  $E_n^*(W_r^*)$  is defined by

$$E_n^*(W_r^*) := \sup_{f \in W_r^*} E_n^*(f).$$

According to Favard [1], and Achieser and Krein [2], we define

$$K_r := (4/\pi) \sum_{m=0}^{\infty} (-1)^{m(r+1)} (2m+1)^{-r-1}. \quad (5)$$

This implies the inequality

$$K_2 < K_4 < \dots < 4/\pi < \dots < K_3 < K_1 = \pi/2. \quad (6)$$

## 2. TURNING THE ALGEBRAIC INTO A TRIGONOMETRIC PROBLEM

The purpose of this section is to replace  $E_n(g)$  by  $E_n^*(g \circ \sin)$ , and to deduce upper bounds for  $E_n(W_1)$  and  $E_n(W_2)$ .

**THEOREM 2.** *Let  $r \in \mathbb{N}$ ,  $n \geq r-1$ ,  $g \in W_r$ , and let*

$$f(t) := \int_0^{\sin t} g^{(r)}(u) (\sin t - u)^{r-1} du / (r-1)! \quad (7)$$

Then

$$E_n(g) = E_n^*(f).$$

*Proof.* Let  $p$  be the best approximation of  $g$  in  $\mathcal{P}_n$ . Then  $g-p$  has at least  $n+2$  alternation points, and  $(g-p) \circ \sin$  has at least  $2n+2$  alternation points in  $[0, 2\pi)$ . Therefore  $p \circ \sin$  is the best approximation of  $g \circ \sin$ , and

$$E_n(g) = \|g-p\| = \|g \circ \sin - p \circ \sin\|^* = E_n^*(g \circ \sin). \quad (8)$$

By Taylor's theorem we have

$$g(\sin t) = \sum_{k=0}^{r-1} g^{(k)}(0) \cdot (\sin t)^k / k! + f(t). \quad (9)$$

Since  $n \geq r-1$ , we have  $E_n^*(g \circ \sin) = E_n^*(f)$ . ■

**THEOREM 3.** *Let  $r \in \mathbb{N}_0$ ,  $f \in C_{2\pi}^r$ ,  $n \in \mathbb{N}_0$ , and let  $\omega(h)$  be a concave modulus of continuity of  $f^{(r)}$ . Then*

$$E_n^*(f) \leq \frac{1}{2} K_r (n+1)^{-r} \omega\left(\frac{\pi}{n+1}\right). \quad (10)$$

*Proof.* See [3].

To prove Theorem 1 for  $r = 1, 2$  we need some special considerations listed in the following theorems:

**THEOREM 4.** *Let  $n \in \mathbb{N}_0$ . Then*

$$E_n(W_1) \leq \sin(\pi/2(n+1)) < K_1/(n+1). \quad (11)$$

*Proof.* Let  $g \in W_1$ ,  $0 \leq h \leq \pi$ ,  $a, b \in \mathbb{R}$ ,  $0 \leq b - a \leq h$ ; for  $f$  see (7).  $\omega(h) := 2 \sin(h/2)$  is a concave modulus of continuity of  $f$ , as

$$|f(b) - f(a)| \leq |\sin b - \sin a| \leq 2 \sin(h/2).$$

By Theorem 3 we get

$$E_n(g) = E_n^*(f) \leq \frac{1}{2} \omega(\pi/(n+1)) = \sin(\pi/2(n+1)). \quad \blacksquare$$

**THEOREM 5.** *Let  $n \in \mathbb{N}$ . Then*

$$E_n(W_2) \leq K_2(n+1)^{-2} < K_2(n-1)!/(n+1)!. \quad (12)$$

*Proof.* Let  $g \in W_r$ ,  $t \in \mathbb{R}$ ; for  $f$  see (7). Then

$$\begin{aligned} |f''(t)| &= \left| (\cos t)^2 g''(\sin t) - \sin t \int_0^{\sin t} g''(u) du \right| \\ &\leq (\cos t)^2 + (\sin t)^2 = 1. \end{aligned} \quad (13)$$

Therefore we obtain that  $f \in W_2^*$ , and by (1) and Theorem 2 we have

$$E_n(g) = E_n^*(f) \leq K_2(n+1)^{-2}. \quad \blacksquare$$

### 3. AN UPPER BOUND OF $E_n(W_r)$

To prove Theorem 1 for  $r \geq 3$ , we need several definitions and lemmas. For all  $r \geq 3$  and  $j \in \mathbb{N}_0$  let

$$p_{rj}(t) := ((d/dt)^{r+j-1} (\sin t - u)^{r-1})_{u=\sin t} / (r-1)!, \quad (14)$$

and

$$B_{rj} := \|\cos \cdot p_{rj}\|^*. \quad (15)$$

It is obvious that  $p_{rj} \in \mathcal{E}_{r-1}$ .

LEMMA 6. Let  $r \geq 3$ ,  $n \geq r - 1$  and  $g \in W_r$ . Then

$$E_n(g) \leq \sum_{j=0}^{\infty} B_{rj} K_{r+j} (n+1)^{-r-j}. \quad (16)$$

*Proof.* Using Theorem 2 we only have to prove the inequality

$$E_n^*(f) \leq \sum_{j=0}^{\infty} B_{rj} K_{r+j} (n+1)^{-r-j}. \quad (17)$$

We split up  $f$  into several functions  $f_j$  so that  $B_{rj}^{-1} \cdot f_j \in W_{r+j}^*$ . For all  $j$  and  $s \in \mathbb{N}_0$  we define  $f_j$  and  $\tilde{f}_s$  by

$$\begin{aligned} f_j &\in C_{2\pi}, & \int_K f_j &= 0, \\ f_j^{(r+j)}(t) &= g^{(r)}(\sin t) \cdot \cos t \cdot p_{rj}(t) + \text{const.}, \end{aligned} \quad (18)$$

and

$$\begin{aligned} \tilde{f}_s &\in C_{2\pi}, & \int_K \tilde{f}_s &= \int_K f, \\ \tilde{f}_s^{(r+s)}(t) &= ((r-1)!)^{-1} \int_0^{\sin t} g^{(r)}(\sin t) \left(\frac{d}{dt}\right)^{r+s} \\ &\quad \cdot (\sin t - u)^{r-1} du + \text{const.} \end{aligned} \quad (19)$$

By calculation of  $f^{(r)}$  we get

$$f^{(r)} = f_0^{(r)} + \tilde{f}_0^{(r)} + \text{const.} \quad (20)$$

Since  $\int_K f^{(r)} = 0 = \int_K f_0^{(r)} + \int_K \tilde{f}_0^{(r)}$  and  $\int_K f = \int_K f_0 + \int_K \tilde{f}_0$  we obtain  $f = f_0 + \tilde{f}_0$ . We also have

$$\tilde{f}_{s-1}^{(r+s)} = (d/dt) \tilde{f}_{s-1}^{(r+s-1)} = f_s^{(r+s)} + \tilde{f}_s^{(r+s)} + \text{const.} \quad (21)$$

By similar arguments as above we obtain  $\tilde{f}_{s-1} = f_s + \tilde{f}_s$ , and thus for all  $s \in \mathbb{N}_0$

$$E_n^*(f) = E_n^* \left( \sum_{j=0}^s f_j + \tilde{f}_s \right) \leq \sum_{j=0}^{\infty} E_n^*(f_j) + E_n^*(\tilde{f}_s). \quad (22)$$

Using Theorem 3 and

$$|f_j^{(r+j)}(t) - f_j^{(r+j)}(t')| \leq 2 \|\cos \cdot p_{rj}\|^* = 2B_{rj}$$

we obtain

$$E_n^*(f_j) \leq B_{rj} K_{r+j} (n+1)^{-r-j}. \quad (23)$$

Since

$$\begin{aligned} & |\tilde{f}_s^{(r+s)}(t) - \tilde{f}_s^{(r+s)}(t')| \\ & \leq 2 \int_0^1 \left\| \left( \frac{d}{dt} \right)^{r+s} (\sin t - u)^{r-1} \right\|^* du / (r-1)! \\ & \leq 2 \int_0^1 (r-1)^{r+s} (1+u)^{r-1} du / (r-1)! \\ & < 2^{r+1} (r-1)^{r+s} / r!, \end{aligned} \quad (24)$$

we have

$$\lim_{s \rightarrow \infty} E_n^*(\tilde{f}_s) \leq \lim_{s \rightarrow \infty} (2^r / r!) K_{r+s} ((r-1)/(n+1))^{r+s} = 0. \quad (25)$$

Combining (22), (23) and (25) we finally get (17). ■

LEMMA 7. Let  $m \geq r \geq 3$ . Then

$$\sum_{j=0}^{\infty} B_{rj} m^{-r-j} \leq (1 - m^{-1}) \cdot (m-r)! / m!. \quad (26)$$

*Proof.* For  $r=3$  we have

$$B_{3j} = \|\cos t \cdot ((\cos 2t)^{(j)} + \sin t \cdot (\sin t)^{(j)})\|^* \leq 2^j$$

and

$$\begin{aligned} \sum_{j=0}^{\infty} B_{3j} m^{-3-j} & \leq m^{-3} \sum_{j=0}^{\infty} (2/m)^j \\ & = (1 - m^{-1}) \cdot (m-3)! / m!. \end{aligned} \quad (27)$$

Now suppose (26) being proved for  $r$ . We prove (26) for  $r+1$ : For  $j \geq 1$  we get  $p_{r+1,j} = \cos \cdot p_{rj} + p'_{r+1,j-1}$ . Since  $p_{r+1,0} = \cos^r = \cos \cdot p_{r0}$ , we obtain

$$p_{r+1,j} = \sum_{k=0}^j (\cos \cdot p_{r,j-k})^{(k)}. \quad (28)$$

Using Bernstein's inequality we have

$$B_{r+1,j} \leq \|p_{r+1,j}\|^* \leq \sum_{k=0}^j r^k B_{r,j-k}, \quad (29)$$

and therefore

$$\begin{aligned} & \sum_{j=0}^{\infty} B_{r+1,j} m^{-r-1-j} \\ & \leq \sum_{j=0}^{\infty} B_{rj} m^{-r-j} \cdot m^{-1} \cdot \sum_{k=0}^{\infty} (r/m)^k \\ & = (m-r)^{-1} \cdot \sum_{j=0}^{\infty} B_{rj} m^{-r-j} \leq (1-m^{-1}) \cdot (m-r-1)!/m!. \quad \blacksquare \end{aligned}$$

LEMMA 8. Let  $m \geq r \geq 3$ . Then

$$\sum_{j=0}^{\infty} B_{rj} K_{r+j} m^{-r-j} < K_r (m-r)!/m!. \quad (30)$$

*Proof.* For  $r$  odd we have  $K_r \geq K_{r+j}$  (see (6)), and by Lemma 7 we get

$$\sum_{j=0}^{\infty} B_{rj} K_{r+j} m^{-r-j} \leq (1-m^{-1}) \cdot K_r \cdot (m-r)!/m!. \quad (31)$$

For  $r$  even we have  $K_{r+1} \geq K_{r+j}$ , and thus

$$\begin{aligned} & \sum_{j=0}^{\infty} B_{rj} K_{r+j} m^{-r-j} \\ & \leq K_r \sum_{j=0}^{\infty} B_{rj} m^{-r-j} + (K_{r+1} - K_r) \sum_{j=1}^{\infty} B_{rj} m^{-r-j} \\ & \leq K_r (1-m^{-1})(m-r)!/m! + (K_{r+1} - K_r)((m-r)!/m! - m^{-r}) \\ & = (1-m^{-1} + (K_{r+1}/K_r - 1)(1-m^{-r}m!/(m-r)!)) K_r (m-r)!/m!. \quad (32) \end{aligned}$$

Now we use two simple estimates, namely,

$$1 - m^{-r} \cdot m!/(m-r)! < 1 - m^{-r}(m-r)^r = 1 - (1-r/m)^r < r^2/m, \quad (33)$$

and

$$\begin{aligned} K_{r+1}/K_r & = \left( \sum_{m=0}^{\infty} (1+2m)^{-r-2} \right) \bigg/ \left( \sum_{m=0}^{\infty} (-1)^m (1+2m)^{-r-1} \right) \\ & < \left( 1 + \sum_{m=1}^{\infty} 3^{-r}(1+2m)^{-2} \right) \bigg/ (1-3^{-r-1}) \\ & = (1 + (\pi^2/8 - 1)3^{-r}) / (1-3^{-r-1}) \\ & = 1 + (\pi^2/8 - 1 + 3^{-1}) / (3^r - 3^{-1}) \\ & < 1 + 1/((1+2)^r - 1) \\ & < 1 + 1/(1+2r+2r(r-1)-1) = 1 + 1/2r^2. \quad (34) \end{aligned}$$

Combining (32), (33) and (34) we obtain

$$\sum_{j=0}^{\infty} B_{rj} K_{r+j} m^{-r-j} < (1 - m^{-1} + 2^{-1} m^{-1}) K_r (m-r)! / m! \quad \blacksquare \quad (35)$$

Now Theorem 1 follows by Lemma 6 and Lemma 8 for  $r \geq 3$ , and by Theorem 4, resp. 5 for  $r = 1$ , resp. 2.

#### REMARKS

Another representation of Theorem 1 is the following.

**COROLLARY 9.** *Let  $r \in \mathbb{N}$ ,  $n \geq r - 1$ ,  $g \in C^r[-1, 1]$ . Then there exists a polynomial  $p \in \mathcal{P}_n$  with*

$$\|g - p\| \leq \|g^{(r)}\| \cdot K_r \cdot (n+1-r)! / (n+1)! \quad (36)$$

*Proof.* If  $\|g^{(r)}\| = 0$ , then  $g \in \mathcal{P}_{r-1} \subset \mathcal{P}_n$ . If  $\|g^{(r)}\| \neq 0$ , (36) follows by Theorem 1 and  $g/\|g^{(r)}\| \in W_r$ .  $\blacksquare$

Note that for small  $n$  the upper bound of  $E_n(W_r)$  given is not very good. For  $n = r - 1$  it is known (see Fisher [5]) that

$$E_{r-1}(W_r) = 2^{1-r}/r!,$$

while our inequality gives only

$$E_{r-1}(W_r) < K_r / r!.$$

#### REFERENCES

1. J. FAVARD, Sur les meilleures procédés d'approximation de certaines classes des fonctions par des polynômes trigonométriques, *Bull. Sci. Math.* **61** (1937), 109-224; 243-256.
2. N. I. ACHESER AND M. G. KREIN, On the best approximation of periodic functions, *Dokl. Akad. Nauk SSSR* **15** (1937), 107-112.
3. N. P. KORNEICHUK, On methods of investigating extremal problems in the theory of best approximation, *Russian Math. Surveys* **29**, No. 3 (1974), 7-43.
4. S. N. BERNSTEIN, "Collected Works," Vol. II, Akad. Nauk SSSR, Moscow, 1954.
5. S. D. FISHER, Best approximation by polynomials, *J. Approx. Theory* **21** (1977), 43-59.