



ELSEVIER

Available at

www.ElsevierMathematics.com

POWERED BY SCIENCE @ DIRECT®

Journal of Approximation Theory 125 (2003) 85–105

JOURNAL OF  
Approximation  
Theory

http://www.elsevier.com/locate/jat

# Multivariate approximating averages

E. Belinsky,<sup>a</sup> F. Dai,<sup>b</sup> and Z. Ditzian<sup>c,\*</sup>

<sup>a</sup>Department of Mathematics, University of the West Indies, Barbados

<sup>b</sup>School of Mathematics, University of Sydney, NSW 2006, Australia

<sup>c</sup>Department of Mathematics, University of Alberta, Edmonton, Alberta, Canada T6G 2G1

## Abstract

Averages on the sphere about  $x$  in  $R^d$  and on the rim of the cap about  $x$  in  $S^{d-1}$  and their iterates are shown to be smoother than  $f$ . Furthermore, their approximating properties satisfy a strong converse inequality of type  $A$  when dealing with multivariate approximation ( $d \geq 2$  in case of  $R^d$  and  $d \geq 3$  in case of  $S^{d-1}$ ). These results are in contrast to the classical results on  $R$  or  $T$  for which the situation is completely different.

© 2003 Published by Elsevier Inc.

MSC: 41A25; 41A63; 42B15

**Keywords:** Moduli of smoothness; Strong converse inequality; Smoothness of an averaging operator; Multipliers

## 1. Introduction

For a function  $f(x)$  on  $R$  or  $T$ , i.e. single variable, the average

$$A_t f(x) = \frac{1}{2}(f(x+t) + f(x-t)) \quad (1.1)$$

approximates  $f(x)$  when  $f(x) \in C(R)(C(T))$  or  $L_p(R)(L_p(T))$ ,  $1 \leq p < \infty$ ; and the rate of approximation relates to the modulus of smoothness  $\omega^2(f, t)_p$  by

$$\sup_{0 < u \leq t} \|f - A_u f\|_p \equiv \omega^2(f, t)_p \approx \inf_g (\|f - g\|_p + t^2 \|g''\|_p). \quad (1.2)$$

\*Corresponding author. Fax: 780-492-6826.

E-mail addresses: ebelinsky@uwichell.edu.bb (E. Belinsky), feng@maths.usyd.edu.au (F. Dai), zditzian@math.ualberta.ca (Z. Ditzian).

Supported by NSERC A4816 of Canada. (Including visits of E. Belinsky and F. Dai in Edmonton.)

It is well-known that  $A_t f$  and its iterates  $A_t^m f$  are not in general smoother than  $f(x)$  and that the supremum on the left of (1.2) cannot be dropped.

We will show that some multivariate analogues which are widely used behave better and the behaviour improves with the dimension.

For  $f(x)$ ,  $x \in \mathbf{R}^d$  (or  $\mathbf{T}^d$ ) we define

$$V_t(f, x) \equiv \frac{1}{m(t)} \int_{\sum_i = \{y: |x-y|=t\}} f(y) d\sigma, \quad V_t(1, x) = 1 \tag{1.3}$$

with  $d\sigma$  being the Lebesgue measure on the sphere  $\sum_i$ .

For  $f(x)$ ,  $x \in S^{d-1} = \{x \in \mathbf{R}^d: |x| = 1\}$ , we define

$$S_t(f, x) = \frac{1}{m(t)} \int_{\gamma_t = \{y: y \cdot x = \cos t\}} f(y) d\gamma, \quad S_t(1, x) = 1, \tag{1.4}$$

where  $d\gamma$  is the Lebesgue measure on the rim of the cap of sphere  $\gamma_t$ .

It turns out that for  $d \geq 2$

$$\|V_t f(\cdot) - f(\cdot)\|_{L_p(\mathbf{R}^d)} \approx \inf_{g \in C^2(\mathbf{R}^d)} (\|f - g\|_{L_p(\mathbf{R}^d)} + t^2 \|\Delta g\|_{L_p(\mathbf{R}^d)}) \tag{1.5}$$

and for  $d \geq 3$

$$\|S_t f(\cdot) - f(\cdot)\|_{L_p(S^{d-1})} \approx \inf_{g \in C^2(S^{d-1})} (\|f - g\|_{L_p(S^{d-1})} + t^2 \|\tilde{\Delta} g\|_{L_p(S^{d-1})}), \tag{1.6}$$

where  $\Delta$  and  $\tilde{\Delta}$  are the Laplacian and the Laplace–Beltrami (the tangential component of the Laplacian on  $S^{d-1}$ ), respectively.

We note that in (1.5) and in (1.6) proved here, there is no supremum on the left-hand side as is common in texts on the subject and as is necessary in the one-dimensional case. Equivalences (1.5) and (1.6) constitute strong converse inequalities of type A in the sense of [Di-Iv], but in one dimension (1.2) is a strong converse inequality of type D in that scale.

We show further that  $V_t^m f$  and  $S_t^m f$  are smoother than  $f$ , and the improvement depends on  $m$ ,  $p$  (of  $L_p$ ) and the dimension  $d$ . In fact, the higher the dimension, the bigger the improvement in smoothness of  $S_t^m f$  and  $V_t^m f$  over that of  $f$ . We note that in one dimension an improvement of smoothness of  $A_t^m f$  is not the case for any  $p$  or  $m$ .

For the proof we will need and show that

$$\|\Delta V_t^m(f)\|_{L_p(\mathbf{R}^d)} \leq \frac{\varepsilon(m)}{t^2} \|f\|_{L_p(\mathbf{R}^d)}, \quad \text{for } d \geq 2 \tag{1.7}$$

and

$$\|\tilde{\Delta} S_t^m f\|_{L_p(S^{d-1})} \leq \frac{\varepsilon(m)}{t^2} \|f\|_{L_p(S^{d-1})} \quad \text{for } d \geq 3, \tag{1.8}$$

where  $\varepsilon(m) = o(1)$ ,  $m \rightarrow \infty$ .

We note that for  $A_t^m$  we cannot guarantee the existence of a derivative, and hence there is no hope for an analogue of (1.7) and (1.8).

The technique we use relies on multipliers. The smoothness of functions on  $S^{d-1}$  will also be related to a new concept introduced recently in [Di-I].

We hope the present results will be helpful in other investigations, though we cannot identify the optimal improvement in smoothness except in the  $L_2$  case.

## 2. Smoothness of $S_t^m f$ in $L_2(S^{d-1})$

We first show the smoothness of  $S_t^m f$  in  $L_2(S^{d-1})$ , which is better than the result which we get for  $L_p$  in the next section (see also Section 7) when we substitute  $p = 2$ . The results in this section will serve as a model and encouragement for later sections and some of the estimates will be used later.

We have (see for instance [Li-Ni,Ru,Wa-Li, (12.4.6), p. 61])

$$S_\theta^m f = \sum_{k=0}^{\infty} \left( \frac{P_k^{(\lambda)}(\cos \theta)}{P_k^{(\lambda)}(1)} \right)^m P_k(f) \equiv \sum_{k=0}^{\infty} (Q_k^{(\lambda)}(\cos \theta))^m P_k(f), \tag{2.1}$$

where  $P_k^{(\lambda)}(x)$  is the ultraspherical (Gegenbauer) polynomial of order  $k$ ,  $\lambda = \frac{d-2}{2}$ ,  $P_k(f)$  is the projection of  $f$  on  $H_k = \{\psi : \tilde{\Delta}\psi = -k(k+d-2)\psi\}$ ,  $\tilde{\Delta}f(x) = \Delta f\left(\frac{x}{|x|}\right)$  (the Laplace–Beltrami operator) and  $\Delta$  is the Laplacian. As both  $S_\theta^m$  and  $\tilde{\Delta}$  are multiplier operators, we have formally

$$-\tilde{\Delta}S_\theta^m f \sim \sum_{k=1}^{\infty} k(k+d-2) \left( \frac{P_k^{(\lambda)}(\cos \theta)}{P_k^{(\lambda)}(1)} \right)^m P_k(f). \tag{2.2}$$

**Theorem 2.1.** For  $f \in L_2(S^{d-1})$ ,  $d \geq 3$ ,  $\theta \leq \frac{\pi}{2}$ , and  $m \geq \frac{4}{d-2}$  we have

$$\|\tilde{\Delta}S_\theta^m f\|_2 \leq \frac{C}{\theta^2} \|f\|_2, \tag{2.3}$$

with  $C$  independent of  $\theta$  and  $f$ .

**Proof.** Using [Sz, (7.33.6)], we have (recalling that there  $P_k^\lambda(1) = \binom{k+2\lambda-1}{k}$ )

$$|Q_k^{(\lambda)}(\cos \theta)| \equiv \left| \frac{P_k^{(\lambda)}(\cos \theta)}{P_k^{(\lambda)}(1)} \right| \leq C_1 \min\{k^{-\lambda}\theta^{-\lambda}, 1\},$$

and hence (recalling  $\lambda = \frac{d-2}{2}$ ) for  $k\theta \geq 1$ ,  $\theta \leq \frac{\pi}{2}$ , we have

$$|\theta^2 k(k+d-2)(Q_k^{(\lambda)}(\cos \theta))^m| \leq C_2 (k\theta)^{2-\frac{d-2}{2}m}$$

which is bounded for  $\frac{d-2}{2}m \geq 2$  or  $m \geq \frac{4}{d-2}$ . For  $k\theta \leq 1$

$$|\theta^2 k(k+d-2)(Q_k^{(\lambda)}(\cos \theta))^m| \leq C_2,$$

which completes the proof.  $\square$

**Corollary 2.2.** For  $f \in L_2(S^{d-1})$ ,  $0 < \theta \leq \frac{\pi}{2}$ ,  $d \geq 3$  and  $m_1 \geq \frac{2}{d-2}$

$$\| \text{grad}_{\tan} S_{\theta}^{m_1} f \|_2 \leq \frac{C}{\theta} \|f\|_2, \tag{2.4}$$

with  $C$  independent of  $\theta$  and  $f$ .

**Proof.** We have for smooth  $f$

$$\begin{aligned} \| \text{grad}_{\tan} S_{\theta}^{m_1} f \|_2^2 &= \langle \text{grad}_{\tan} S_{\theta}^{m_1} f, \text{grad}_{\tan} S_{\theta}^{m_1} f \rangle \\ &= \langle -\tilde{\Delta} S_{\theta}^{2m_1} f, f \rangle \leq \| \tilde{\Delta} S_{\theta}^{2m_1} f \|_2 \|f\|_2 \\ &\leq \frac{C}{\theta^2} \|f\|_2 \|f\|_2, \end{aligned}$$

and as the last result is independent of the smoothness of  $f$ , (2.4) follows.  $\square$

We can also prove

**Theorem 2.3.** For  $f \in L_2(S^{d-1})$ ,  $d \geq 3$ ,  $0 < \theta \leq \frac{\pi}{2}$ , and  $m$  big enough

$$\| \tilde{\Delta} S_{\theta}^m f \|_2 \leq \frac{\varepsilon(m)}{\theta^2} \|f\|_2, \tag{2.5}$$

where  $\varepsilon(m) = o(1)$  as  $m \rightarrow \infty$ .

We need the following lemma, which will also be used later.

**Lemma 2.4.** For  $n\theta \leq 1$ ,  $n \geq 1$ ,  $0 < \theta \leq \frac{\pi}{2}$  and  $\lambda = \frac{d-2}{2}$

$$\left| \frac{P_n^{(\lambda)}(\cos \theta)}{P_n^{(\lambda)}(1)} \right| \equiv |Q_n^{(\lambda)}(\cos \theta)| \leq 1 - \frac{n^2}{d-1} \sin^2 \frac{\theta}{2} \leq 1 - \frac{2}{\pi^2(d-1)} n^2 \theta^2. \tag{2.6}$$

For  $n\theta \geq 1$ ,  $n \geq 1$ ,  $0 < \theta \leq \frac{\pi}{2}$  and  $\lambda = \frac{d-2}{2}$

$$|Q_n^{(\lambda)}(\cos \theta)| < \alpha < 1 \tag{2.7}$$

with  $\alpha$  independent of  $n$  and  $\theta$  (but depending on the fact that  $n\theta \geq 1$  and on  $\lambda$ ).

**Proof.** To show (2.6) we integrate by parts and use [Sz, (4.7.14), p. 82] with  $\lambda = \frac{d-2}{2}$  to obtain for  $n \geq 1$

$$\begin{aligned} 1 - Q_n^{(\lambda)}(\cos t) &= - \int_0^{\theta} \frac{d}{du} Q_n^{(\lambda)}(\cos u) du \\ &= \frac{n(n+d-2)}{d-1} \int_0^{\theta} Q_{n-1}^{(\lambda+1)}(\cos u) \sin u du \\ &= \frac{n(n+d-2)}{d-1} \left[ \int_0^{\theta} \sin u du - \int_0^{\theta} \left( 1 - Q_{n-1}^{(\lambda+1)}(\cos u) \right) \sin u du \right]. \end{aligned}$$

For  $n = 1$   $Q_n^{(\lambda)}(\cos \theta) = \cos \theta = 1 - 2 \sin^2 \frac{\theta}{2}$ . Otherwise, we use the mean value theorem to obtain

$$\begin{aligned} 0 \leq 1 - Q_{n-1}^{(\lambda+1)}(\cos u) &= (1 - \cos u) \frac{(n-1)(n+d-1)}{d+1} Q_{n-2}^{(\lambda+2)}(\cos \xi) \\ &\leq (1 - \cos u) \frac{(n-1)(n+d-1)}{d+1}, \end{aligned}$$

and hence

$$\int_0^\theta \left(1 - Q_{n-1}^{(\lambda+1)}(\cos u)\right) \sin u \, du \leq \frac{(n-1)(n+d-1)}{d+1} \frac{(1 - \cos \theta)^2}{2}.$$

For  $n\theta < 1$  we have  $\frac{(n-1)(n+d-1)}{d+1} (1 - \cos \theta) \leq 1$ , and hence

$$\begin{aligned} 0 \leq 1 - 2 \frac{n(n+d-2)}{d-1} \sin^2 \frac{\theta}{2} &\leq Q_n^{(\lambda)}(\cos \theta) \\ &\leq 1 - \frac{n(n+d-2)}{d-1} \sin^2 \frac{\theta}{2} \\ &\leq 1 - \frac{n^2}{d-1} \sin^2 \frac{\theta}{2} \leq 1 - \frac{2}{(d-1)\pi^2} (n\theta)^2. \end{aligned}$$

For  $n\theta \geq 1$ , we use the identity [Sz, (4.10.3), p. 97]

$$\begin{aligned} Q_n^{(\lambda)}(\cos \theta) &= \left( \int_0^\pi \sin^{2\lambda-1} \varphi \, d\varphi \right)^{-1} \int_0^\pi (\cos \theta + i \sin \theta \cos \varphi)^n \sin^{2\lambda-1} \varphi \, d\varphi \\ &= \left( \int_0^\pi \sin^{2\lambda-1} \varphi \, d\varphi \right)^{-1} \int_0^\pi \operatorname{Re}(\cos \theta + i \sin \theta \cos \varphi)^n \sin^{2\lambda-1} \varphi \, d\varphi. \end{aligned} \tag{2.8}$$

For  $\theta_0 \leq \theta \leq \pi - \theta_0$  we find  $\delta$  such that

$$\int_0^\delta \sin^{2\lambda-1} \varphi \, d\varphi \Big/ \int_0^\pi \sin^{2\lambda-1} \varphi \, d\varphi = \frac{1}{8},$$

and estimate for  $n \geq 2$

$$\begin{aligned} |Q_n^{(\lambda)}(\cos \theta)| &\leq \frac{1}{4} + \frac{3}{4} |\cos \theta + i \sin \theta \cos \delta|^n \\ &= \frac{1}{4} + \frac{3}{4} (1 - \sin^2 \theta \sin^2 \delta)^{n/2} \\ &\leq \frac{1}{4} + \frac{3}{4} (1 - \sin^2 \theta \sin^2 \delta) \\ &\leq 1 - \frac{3}{4} \sin^2 \theta_0 \sin^2 \delta, \end{aligned}$$

and for  $n = 1$   $Q_n^{(\lambda)}(\cos \theta) = \cos \theta \leq \cos \theta_0 = 1 - 2 \sin^2 \frac{\theta_0}{2}$ , and hence (2.7) is valid for  $\theta_0 \leq \theta < \pi - \theta_0$  (not only for  $\theta < \frac{\pi}{2}$ ) and  $\alpha$  depending on  $\theta_0$  and  $\lambda$  (but not yet on  $n\theta$ ). We choose  $\theta_0$  so that  $\cos \theta_0 \geq 0.9$  (for example) and note that Fejer showed [Sz, (6.6.6), p. 138] that the zeros of  $P_n^{(\lambda)}(\cos \theta)$  satisfy

$$\frac{v - (1 - \lambda)/2}{n + \lambda} \pi < \theta_v < \frac{v + \lambda - \frac{1}{2}}{n + 2\lambda} \pi,$$

and hence the next extremum of  $P_n^{(\lambda)}(\cos \theta)$  after  $\theta = 0$  is at a zero of  $P_{n-1}^{(\lambda+1)}(\cos \theta)$  i.e.

$$\theta \leq \frac{1+\lambda+\frac{1}{2}}{n-1+2\lambda+2} \pi = \frac{1}{2} \frac{d+1}{n+d-1} \pi \leq \frac{1}{2} \frac{d+1}{n} \pi.$$

As the  $|P_n^{(\lambda)}(\cos \theta)|$  at the extrema are descending for  $0 \leq \theta \leq \frac{\pi}{2}$  (increasing in  $x = \cos \theta$ ) [Sz, (1), pp. 168–169], we have to check the values only in the range  $1 \leq \theta n \leq \frac{(d+1)\pi}{2}$ . Using (2.8), we note that for  $n \leq \frac{(d+1)\pi}{2\theta}$  and  $\tan^{-1} \frac{\sin \theta \cos \varphi}{\cos \theta} < \frac{\pi}{2n}$ ,  $|\arg(\cos \theta + i \sin \theta \cos \varphi)^n| < \frac{\pi}{2}$ , and hence  $\operatorname{Re}(\cos \theta + i \sin \theta \cos \varphi)^n > 0$ . Clearly,  $\tan^{-1} \left( \frac{\sin \theta \cos \varphi}{0.9} \right) < \tan^{-1} \frac{\theta \cos \varphi}{0.9} < \frac{\theta \cos \varphi}{0.9}$  and hence for  $\frac{\theta \cos \varphi}{0.9} < \frac{\pi}{2n}$  or  $\cos \varphi \leq 0.9 \frac{\pi}{2n\theta} \leq 0.9 \frac{\pi}{(d+1)\pi} = 0.9 \frac{1}{d+1}$ . We choose  $\varphi_0$  such that  $\cos \varphi_0 = 0.9 \frac{1}{d+1}$  and as

$$\left( \int_{\varphi_0}^{\pi-\varphi_0} \sin^{2\lambda-1} \varphi \, d\varphi \right) / \int_0^\pi \sin^{2\lambda-1} \varphi \, d\varphi \geq \beta > 0.$$

with  $\beta$  depending on  $\lambda$  only, we have for  $\theta n \leq \frac{(d+1)\pi}{2}$ , using (2.8)

$$Q_n^{(\lambda)}(\cos \theta) \geq -(1 - \beta) \max |\cos \theta + i \sin \theta \sin \varphi|^n \geq -1 + \beta,$$

and hence  $Q_n^{(\lambda)}(\cos \theta) \geq -1 + \beta$  at the first extremum after  $\theta = 0$ . Therefore, for  $n\theta \geq 1$ ,  $\theta \leq \theta_0 < \frac{\pi}{2}$

$$|Q_n^{(\lambda)}(\cos \theta)| \leq \max \left( 1 - \frac{2}{\pi^2} \frac{1}{d-1}, 1 - \beta \right) = \alpha < 1. \quad \square$$

**Proof of Theorem 2.3.** For  $n\theta \leq 1$  we use (2.6) of Lemma 2.4 to estimate the coefficients

$$\begin{aligned} \theta^2 n(n+d-2) \left( 1 - \frac{2}{\pi^2(d-1)} (n\theta)^2 \right)^m &\leq (d-1)\theta^2 n^2 \left( 1 - \frac{2}{\pi^2(d-1)} (n\theta)^2 \right)^m \\ &\leq C \frac{1}{m} = o(1), \quad m \rightarrow \infty. \end{aligned}$$

For  $n\theta \geq 1$  we write  $m = m_1 + m_2$  where  $m_1 \geq \frac{4}{d-2}$  from Theorem 2.1, and use (2.7) of Lemma 2.4 to get  $\alpha^{m_2} = o(1)$ ,  $m_2 \rightarrow \infty$ , which concludes the proof.  $\square$

### 3. Smoothness of $S_\theta^m f$ for $f \in L_p(S^{d-1})$ , $1 \leq p \leq \infty$

In this section we exhibit the smoothness of  $S_\theta^m f$  for some power  $m$ .

**Theorem 3.1.** For  $f \in L_p(S^{d-1})$ ,  $1 \leq p \leq \infty$ ,  $d \geq 3$  and  $m > \frac{2(\binom{d}{2}+3)}{d-2}$  we have

$$\|\tilde{\Delta} S_\theta^m f\|_p \leq C \max \left( \frac{1}{\theta^2}, \frac{1}{(\pi - \theta)^2} \right) \|f\|_p. \tag{3.1}$$

**Remark 3.2.** Theorem 3.1 does not contain Theorem 2.1 as the condition on  $m$  is harsher. In Section 7 we will improve the condition on  $m$  to be close to that in Theorem 2.1 for  $p$  close to 2.

We need the following lemma:

**Lemma 3.2.** For integers  $j, k$  and  $m$  and for  $0 < \theta \leq \frac{\pi}{2}$  we have

$$\left| \Delta^j \left( Q_k^{(\lambda)}(\cos \theta) \right)^m \right| \leq \begin{cases} C \theta^j / (k\theta)^{m\lambda} & \text{for } k\theta \geq 1, \\ C \theta^j & \text{for } k\theta \leq 1, \end{cases} \tag{3.2}$$

where  $C$  is independent of  $k$  and  $\theta$  but may depend on  $j, \lambda$  and  $m$ , and where  $\Delta a_k \equiv \bar{\Delta} a_k \equiv a_{k+1} - a_k, \Delta^j a_k \equiv \Delta(\Delta^{j-1} a_k)$ .

**Proof.** Using Leibnitz’s theorem for differences, we only have to show (3.2) for  $m = 1$  and all  $j$ . This result follows from classical estimates and was essentially demonstrated in [Da-Wa-Yu, (21)]. For  $j = 0, m = 1$  (3.2) is in [Sz, (7.33.6), p. 170]. For  $\alpha, \beta$  one uses [Sz, (4.5.4), p. 71] to obtain for  $Q_k^{(\alpha, \beta)}(x) \equiv P_k^{(\alpha, \beta)}(x) / P_k^{(\alpha, \beta)}(1)$

$$Q_{k+1}^{(\alpha, \beta)}(x) - Q_k^{(\alpha, \beta)}(x) = -(1-x) \frac{(2k + \alpha + \beta + 2)}{2(\alpha + 1)} Q_k^{(\alpha+1, \beta)}(x)$$

and by induction

$$\Delta^j Q_k^{(\alpha, \beta)}(x) = \sum_{\ell \geq j/2}^j (1-x)^\ell P_\ell(k) Q_k^{(\alpha+\ell, \beta)}(x),$$

where  $P_\ell(k)$  are polynomials of degree  $2\ell - j$  in  $k$ . From the above, using  $\alpha = \lambda - \frac{1}{2}$  with [Sz, (8.21.18), p. 196] for  $k\theta \geq 1$  and  $|Q_k^{(\alpha+j, \beta)}(x)| \leq 1$  for  $k\theta < 1$ , we complete the proof of (3.2).  $\square$

**Remark 3.3.** For  $\theta \geq \frac{\pi}{2}$

$$S_\theta f(x) = S_{\pi-\theta} f(x_0)$$

with  $x_0$  the antipodal of  $x$ , that is the other point of intersection of  $S^{d-1}$  with the line connecting  $x$  with the center. Observe that

$$S_\theta^m f(x) = S_{\pi-\theta}^m f(x) \quad \text{for even } m$$

and

$$S_\theta^m f(x) = S_{\pi-\theta}^m f(x_0) \quad \text{for odd } m.$$

**Proof of Theorem 3.1.** In view of Remark 3.3 we have to prove our result only for  $0 < \theta \leq \frac{\pi}{2}$ . It was shown in [Bo-Cl] that the Cesàro summability of order  $r$  satisfies  $\|\sigma_n^r f\|_{L_p(S^d)} \leq C \|f\|_{L_p(S^d)}$ , for any  $r > \frac{d-2}{2}$  (and actually for  $r > (d-2)|\frac{1}{2} - \frac{1}{p}|$ ). We will

use the well-known result relating

$$-\tilde{\Delta} S_\theta^m f \sim \sum_{k=1}^\infty k(k+d-2) (Q_k^{(\lambda)}(\cos \theta))^m P_k(f) \tag{3.3}$$

to

$$-\theta^2 \tilde{\Delta} S_\theta^m f = \theta^2 \sum_{k=1}^\infty \Delta^{r+1} \left\{ k(k+d-2) (Q_k^{(\lambda)}(\cos \theta))^m \right\} \binom{k+r}{r} \sigma_k^r(f)$$

(which follows from the identity  $P_k(f) = \overleftarrow{\Delta}^{r+1} \binom{k+r}{r} \sigma_k^r(f, x)$ , using summation by parts) if the latter is convergent termwise in norm. It is therefore sufficient to show that

$$\theta^2 \sum_{k=1}^\infty \left| \Delta^{r+1} \left\{ k(k+d-2) (Q_k^{(\lambda)}(\cos \theta))^m \right\} \right| \binom{k+r}{r} \leq C_1$$

or

$$S = \theta^2 \sum_{k=1}^\infty \left| \Delta^{r+1-i} (Q_k^{(\lambda)}(\cos \theta))^m \right| k^{r+2-i} \leq C_2$$

for  $i = 0, 1, 2$ . We separate the sum  $S$  by  $S = S_1 + S_2$  where  $S_1$  sums on  $k$  satisfying  $k\theta \geq 1$  and  $S_2$  when  $k\theta \leq 1$ .

Using Lemma 3.2, we have

$$\begin{aligned} S_1 &\leq C_3 \theta^2 \sum_{k\theta \geq 1} \frac{1}{(k\theta)^{\lambda m}} \theta^{r+1-i} k^{r+2-i} \\ &\leq C_4 \theta^{2-\lambda m+r+1-i} \sum_{k\theta \geq 1} k^{r+2-i-\lambda m}. \end{aligned}$$

When  $r + 2 - i - \lambda m < -1$ , that is,  $\frac{r+3}{\lambda} < m$  or  $\frac{2(r+3)}{d-2} < m$ ,  $S_1$  is bounded.

To estimate  $S_2$  we write

$$S_2 \leq C_5 \theta^2 \sum_{k\theta \leq 1} \left(\frac{2}{k}\right)^{r+1-i} k^{r+2-i} \leq C_6 \theta^2 \sum_{k\theta \leq 1} k \leq C_6. \quad \square$$

Perhaps as an intuitive incentive to reduce  $m$  (but certainly not a proof) we observe that the same method used to prove Theorem 3.1 yields:

**Theorem 3.4.** *For  $f \in L_p(S^{d-1})$   $d \geq 3$  and  $m > \frac{2(r+2\ell+1)}{(d-2)\ell}$  we have*

$$\|\tilde{\Delta}^\ell S_\theta^{\ell m} f\|_p \leq C \max\left(\frac{1}{\theta^{2\ell}}, \frac{1}{(\pi-\theta)^{2\ell}}\right) \|f\|_p. \tag{3.4}$$

We will not elaborate on the proof of Theorem 3.4 as it follows almost word for word that of Theorem 3.1.



**Remark 3.5.** For  $1 < p < \infty$  and  $f \in L_p(S^{d-1})$  we could have now deduced for  $0 < \theta \leq \frac{\pi}{2}$

$$\|\tilde{\Delta} S_\theta^m f\|_p \leq \frac{\varepsilon(m)}{\theta^2} \|f\|_p$$

using Theorems 2.1, 3.1 and the Riesz–Thorin interpolation theorem. However, the result is valid for  $p = 1$  and  $\infty$  as well, as will be shown (with substantial additional work) in Section 4.

#### 4. Estimate of $S_\theta^m f$ for high power $m$

The result of this section is the following theorem.

**Theorem 4.1.** For  $f \in L_p(S^{d-1})$ ,  $1 \leq p \leq \infty$ ,  $d \geq 3$  and  $m$  big enough

$$\|\tilde{\Delta} S_\theta^m f\|_p \leq \varepsilon(m) \max\left(\frac{1}{\theta^2}, \frac{1}{(\pi - \theta)^2}\right) \|f\|_p, \tag{4.1}$$

where  $\varepsilon(m) \rightarrow 0$  as  $m \rightarrow \infty$ .

**Proof.** It is sufficient in view of Remark 3.3 to show (4.1) for  $0 < \theta \leq \frac{\pi}{2}$ . We write, following the proof of Theorem 3.1,

$$-\theta^2 \tilde{\Delta} S_\theta^m f = \theta^2 \sum_{k=1}^\infty \Delta^{r+1} \left( (k(k+d-2)) \left( Q_k^{(\lambda)}(\cos \theta) \right)^m \right) \binom{k+r}{r} \sigma_k^r(f),$$

and we have to show

$$S = \theta^2 \sum_{k=1}^\infty \left| \Delta^{r+1} (k(k+d-2)) \left( Q_k^{(\lambda)}(\cos \theta) \right)^m \right| k^r \leq \varepsilon(m), \quad m \rightarrow \infty.$$

We set

$$S \leq C(d) \theta^2 \sum_{i=0}^2 \sum_{k=1}^\infty \left| \Delta^{r+1-i} Q_k^{(\lambda)}(\cos \theta)^m \right| k^{r+2-i} \equiv \sum_{i=0}^2 S(i).$$

We estimate  $S(0)$ , and estimates for  $S(1)$  and  $S(2)$  are almost identical. We write now, choosing  $\ell > \frac{r+3}{\lambda} \geq \frac{2(d+5)}{d-2}$  and  $\ell \geq r+1$  say  $\ell = \max\left(\left[\frac{2(d+5)}{d-2}\right] + 1, \left[\frac{d-2}{2}\right] + 2\right)$  (which is perhaps lavish),

$$S(0) \leq C(d) \theta^2 \left[ \sum_{1 \leq k \leq \frac{1}{\theta}} + \sum_{k > \frac{1}{\theta}} \right] \left\{ m^{r+1} A_{k,\ell}(\theta) \left( \max_{0 \leq s \leq r+1} \left| Q_{k+s}^{(\lambda)}(\cos \theta) \right| \right)^{m-\ell} k^{r+2} \right\} \\ \equiv S_1 + S_2,$$

where

$$A_{k,\ell}(\theta) = \max \left\{ \prod_{1 \leq i \leq \ell} \left| \Delta^{j_i} Q_{k+s}^{(\lambda)}(\cos \theta) \right|; 0 \leq j_i \leq r+1, \sum_{i=1}^{\ell} j_i = r+1, 0 \leq s \leq r \right\}.$$

For  $S_1$  we note that, using the estimate in Lemma 3.2, we have

$$\begin{aligned} S_1 &\leq C\theta^2 \sum_{1 \leq k \leq \frac{1}{\theta}} m^{r+1} \theta^{r+1} (B_k(\theta))^{m-\ell} k^{r+2} \\ &\leq C\theta^2 \sum_{1 \leq k \leq \frac{1}{\theta}} km^{r+1} (k\theta)^{r+1} (B_k(\theta))^{m-\ell}, \end{aligned}$$

where  $B_k(\theta) = \max_{0 \leq s \leq r+1} |Q_{k+s}^{(\lambda)}(\cos \theta)|$ . Using (2.6) of Lemma 2.4, we have for  $m > 2\ell$  (we note that using (2.7) the range in which (2.6) is valid can be extended to include  $k+s \leq k+r+1$ )

$$m^{r+1} (k\theta)^{r+2} (B_k(\theta))^{m-\ell} \approx \frac{1}{m},$$

and hence

$$S_1 \approx \frac{1}{m} \theta \sum_{1 \leq k \leq \frac{1}{\theta}} \approx \frac{1}{m}.$$

To estimate  $S_2$  we use the estimate in Lemma 3.2 and write

$$S_2 \leq C\theta^2 \sum_{k > \frac{1}{\theta}} [m^{r+1} B_k(\theta)^{m-\ell}] k^{r+2} \frac{\theta^{r+1}}{(k\theta)^{\lambda \ell}}.$$

Using (2.7) of Lemma 2.4,

$$m^{r+1} Q_{k+s}^{(\lambda)}(\cos \theta)^{m-\ell} = o(1) \text{ as } m \rightarrow \infty \text{ and } k\theta > 1 \text{ and } s \leq r+1$$

and hence for  $\ell$  chosen earlier  $S_2 = o(1)$ ,  $m \rightarrow \infty$  in a manner that is dependent only on  $\lambda$  ( $r$  is fixed by  $\lambda$  as well).  $\square$

### 5. The equivalence result and other corollaries

As a corollary of the results in Sections 3 and 4 we will obtain the equivalence result or strong converse inequality of type A (in the terminology of [Di-Iv]).

**Theorem 5.1.** *For  $f \in L_p(S^{d-1})$ ,  $1 \leq p \leq \infty$ ,  $d \geq 3$  and  $0 < \theta \leq \frac{\pi}{2}$*

$$\|S_\theta f - f\|_p \approx \inf_g (\|f - g\|_p + \theta^2 \|\tilde{\Delta}g\|_p) \equiv K(f, \theta^2)_p, \tag{5.1}$$

where the infimum is taken on sufficiently smooth  $g$ .

**Remark 5.2.** The common form of the relationship between  $S_\theta f - f$  and  $K(f, \theta^2)_p$  is

$$\sup_{\theta \leq t} \|S_\theta f - f\|_p \approx \inf_g (\|f - g\|_p + t^2 \|\tilde{\Delta}g\|_p), \tag{5.2}$$

and it is the dropping of the supremum on the left of (5.2) which defined smoothness in all previous papers on the subject (scores of them) that is a highlight in this paper.

To prove (5.1) we follow the method of [Di-Iv, Section 4] but for that we need in addition to Theorem 4.1 the following improved Voronovskaja-type result for  $S_\theta$ .

**Lemma 5.3.** For  $g$  such that  $\tilde{\Delta}^i g \in L_p(S^{d-1})$ ,  $1 \leq p < \infty$ , for  $i = 0, 1, 2$  and  $\tilde{\Delta}^i g \in C$  for  $p = \infty$  and for  $\theta$  satisfying  $0 < \theta \leq \frac{\pi}{2}$

$$\|S_\theta g - g - \alpha(\theta)\tilde{\Delta}g\|_p \leq C\theta^4 \|\tilde{\Delta}^2 g\|_p \tag{5.3}$$

with  $0 < A\theta^2 \leq \alpha(\theta) \leq B\theta^2$  and  $A, B, C$  independent of  $\theta$  and  $g$ .

**Proof.** Using Theorem 2.1 of [Di-RuII] for  $g \in C^2(S^{d-1})$ ,

$$S_\theta(g, x) - g(x) = \frac{\Gamma(\frac{d-1}{2})}{2\pi^{(d-1)/2}} \int_0^\theta \frac{dt}{(\sin t)^{d-2}} \int_{\sigma_x(t)} \tilde{\Delta}g(y) d\sigma(y), \tag{5.4}$$

where  $\sigma_x(t) = \{y; |y| = 1, \cos t \leq x \cdot y \leq 1\}$ . Repeating this process as was done in Lemma 4.2 of [Di-RuII], we obtain (5.3) for  $g \in C^4(S^{d-1})$ . Using density of  $C^4(S^{d-1})$  in the space for which  $\tilde{\Delta}^i g$  for  $i = 0, 1, 2$  are in  $L_p$ ,  $1 \leq p < \infty$  and in  $C$  for  $p = \infty$ , we complete the proof.  $\square$

**Proof of Theorem 5.1.** We follow [Di-Iv, Section 4]. For  $g$  satisfying  $g, \tilde{\Delta}g \in L_p$ , or  $g, \tilde{\Delta}g \in C$  (or  $p = \infty$ ) (5.4) implies

$$\|S_\theta g - g\|_p \leq C\theta^2 \|\tilde{\Delta}g\|_p,$$

and as  $S_\theta$  is a contraction on  $L_p$ ,  $1 \leq p \leq \infty$ , we have the known direct result

$$\|S_\theta f - f\|_p \leq CK(f, \theta^2)_p.$$

To prove the converse (strong converse inequality of type A in the terminology of [Di-Iv]), we use for  $g$  of (5.3)  $S_\theta^m f$  with  $m$  to be chosen. Clearly,

$$\|f - g\|_p = \|f - S_\theta^m f\|_p \leq m \|f - S_\theta f\|_p.$$

As we established that  $S_\theta^m f$  and  $\tilde{\Delta}S_\theta^m f$  are in  $L_p$  for  $m$  big enough, it remains to show that for some  $m$

$$\theta^2 \|\tilde{\Delta}S_\theta^m f\|_p \leq C \|f - S_\theta f\|_p. \tag{5.5}$$

To prove (5.5) we first observe that using Theorem 3.1 with  $m > \frac{4(\lfloor \frac{d}{2} \rfloor + 3)}{d-2}$ ,  $\tilde{\Delta}S_\theta^m f$  and  $\tilde{\Delta}^2 S_\theta^m f$  are in  $L_p$   $1 \leq p \leq \infty$  if  $f$  is. Furthermore, if  $m > \frac{6(\lfloor \frac{d}{2} \rfloor + 3)}{d-2}$  and  $f \in L_\infty(S^{d-1})$ ,

$\tilde{\Delta}^3 f \in L_\infty(S^{d-1})$ , and this implies that  $\tilde{\Delta}^2 f$  is continuous. We now choose  $m = \frac{4(\lfloor \frac{d}{2} \rfloor + 3)}{d-2} + m_1$  where  $m_1 \left( m_1 \geq \frac{2\lfloor \frac{d}{2} \rfloor + 3}{d-2} \right)$  is such that for all  $f \in L_p$

$$\|\tilde{\Delta} S_\theta^{m_1} f\|_p \leq \frac{A}{2C} \frac{1}{\theta^2} \|f\|_p \tag{5.6}$$

with  $A$  and  $C$  of Lemma 5.3. The possibility of such a choice is guaranteed by Theorem 4.1. We use Lemma 5.3 on  $g = S_\theta^m f$  and obtain

$$\begin{aligned} \|\alpha(\theta)\tilde{\Delta} S_\theta^m f\|_p &\leq \|S_\theta^m f - f\|_p + C\theta^4 \|\tilde{\Delta}^2 S_\theta^m f\| \\ &\leq m\|S_\theta f - f\|_p + \frac{A}{2}\theta^2 \|\tilde{\Delta} S_\theta^{m-m_1} f\|_p \\ &\leq m\|S_\theta f - f\|_p + \frac{A}{2}\theta^2 \|\tilde{\Delta} S_\theta^m f\|_p + \frac{A\theta^2}{2} \|\tilde{\Delta} S_\theta^{m-m_1} (S_\theta^{m_1} - I)f\|_p \\ &\leq m\|S_\theta f - f\|_p + \frac{A}{2} C_1 \|(S_\theta^{m_1} - I)f\|_p + \frac{A}{2}\theta^2 \|\tilde{\Delta} S_\theta^m f\|_p, \end{aligned}$$

with  $C_1$ , the constant from Theorem 3.1. Use of the triangle inequality will complete the proof.  $\square$

We also have as a corollary of the above and Theorem 10.4 of [Di-Iv]:

**Corollary 5.4.** *For  $f \in L_p(S^{d-1})$ ,  $1 \leq p \leq \infty$ ,  $d \geq 3$ ,  $0 < \theta \leq \frac{\pi}{2}$  and integer  $\ell$*

$$\|(S_\theta - I)^\ell f\|_p \approx \inf(\|f - g\|_p + \theta^{2\ell} \|\tilde{\Delta}^\ell g\|_p) \equiv K_r(f, \tilde{\Delta}, \theta^{2r})_p. \tag{5.7}$$

**Proof.** We just have to verify that  $S_\theta$  satisfies the conditions (for  $Q_d$ ) in Theorem 10.4 of [Di-Iv], which it does.  $\square$

**6. The result on  $R^d$  or  $T^d$**

We define  $V_t f$  for  $f$  on  $R^d$  or  $T^d$  by

$$V_t(f, x) = \frac{1}{|S^{d-1}|} \int_{S^{d-1}} f(x + ty) d\sigma(y), \tag{6.1}$$

where  $S^{d-1}$  is the unit sphere in  $R^d$ ,  $|S^{d-1}|$  is its measure, and the integration  $d\sigma(y)$  is on the sphere. It is well-known that  $V_t$  is a contraction on  $L_p(R^d)$  or  $L_p(T^d)$ ,  $1 \leq p \leq \infty$ . We will deal with  $R^d$  and show how to copy the results to  $T^d$ .

Using [St-We, p. 154], we have for  $x \in R^d$  and  $v = \frac{d-2}{2}$

$$\widehat{\Delta V_t^m} f(x) = (-4\pi^2|x|^2) \cdot \left( \Gamma\left(\frac{d}{2}\right) \frac{J_v(2\pi t|x|)}{(\pi t|x|)^v} \right)^m \hat{f}(x), \tag{6.2}$$

where

$$\hat{f}(x) = \int_{\mathbf{R}^d} f(y)e^{-2\pi ixy} dy.$$

We can now prove the following result about the smoothness of  $V_t f$ .

**Theorem 6.1.** For  $f \in L_2(\mathbf{R}^d)$ ,  $d \geq 2$  and  $m \geq \frac{4}{d-1}$

$$\|\Delta V_t^m f\|_2 \leq \frac{C}{t^2} \|f\|_2. \tag{6.3}$$

**Proof.** Using (6.2), it is sufficient to show that for  $v = \frac{d-2}{2}$

$$t^2|x|^2 \left| \frac{\Gamma\left(\frac{d}{2}\right) J_v(2\pi t|x|)}{(\pi t|x|)^v} \right|^m = I(m, v, t|x|)$$

is bounded. For  $t|x| \leq 1$   $I(m, v, t|x|)$  is bounded by 1 which follows from the definition of  $J_v(u)$  [St-We, p. 154], as

$$\frac{\Gamma\left(\frac{d}{2}\right)}{\Gamma\left(\frac{d-1}{2}\right)\Gamma\left(\frac{1}{2}\right)} \int_{-1}^1 (1-s^2)^{(d-3)/2} ds = 1.$$

For  $t|x| \geq 1$  we use Lemma 3.11 in [St-We, p. 158] to obtain

$$\frac{|J_v(2\pi t|x|)|}{(\pi t|x|)^v} \leq \frac{C}{(2\pi t|x|)^{v+\frac{1}{2}}}$$

and  $I(m, v, t|x|)$  is bounded if  $2 - m(v + \frac{1}{2}) \leq 0$  or  $m \geq \frac{4}{d-1}$ .  $\square$

**Theorem 6.2.** For  $f \in L_p(\mathbf{R}^d)$ ,  $d \geq 2$  and  $m > \frac{2(d+2)}{d-1}$

$$\|\Delta V_t^m f\|_p \leq \frac{C}{t^2} \|f\|_p. \tag{6.4}$$

**Proof.** A simple change of variables implies that it is sufficient to prove (6.4) for  $t = 1$ . Furthermore, it is sufficient to prove (6.4) for  $p = 1$ , which implies (6.4) for  $p = \infty$  by duality, and for  $1 < p < \infty$  by the Riesz–Thorin Theorem. To prove it now (for  $L_1(\mathbf{R}^d)$ ) it is sufficient to show that

$$|x|^2 \left( \frac{\Gamma\left(\frac{d}{2}\right) J_v(2\pi|x|)}{(\pi|x|)^v} \right)^m \equiv I(m, v, |x|)$$

is in  $L_1(\mathbf{R}^d)$  and is a Fourier transform of an element in  $L_1$  for the given  $m$ . For any  $m$  and  $|x| \leq 1$ ,  $I(m, v, |x|) \leq 1$ . To show that  $I(m, v, |x|)$  is in  $L_1$  it suffices to show for  $|x| \geq 1$   $|I(m, v, |x|)| \leq \frac{C}{|x|^{d+\alpha}}$  with  $\alpha > 0$ . Following Lemma 3.11 of [St-We, p. 158], this estimate is achieved for  $d + 2 < m(\frac{d-1}{2})$  or for  $m > \frac{2(d+2)}{d-1}$ . Therefore,  $I(m, v, |x|)$  is the Fourier transform of a bounded continuous function. To show that it is a Fourier transform of a  $L_1$  function it is sufficient to show that  $d + 1$  derivatives of  $I(m, v, |x|)$  are bounded in  $L_1$ . Using the recursion relation

$$\frac{d}{du}(u^{-v}J_v(u)) = -u^{-v}J_{v+1}(u)$$

(see [St-We, p. 153]) and the split used above, we obtain that the derivatives of  $I(m, v, |x|)$  are in  $L_1$  and hence its inverse Fourier transform is in  $L_1$ . (Actually, it is continuous and of support in  $|u| \leq m$ .) We have  $\|\mathcal{F}^{-1}(I(m, v, |x|))\|_1 \leq C$ , and hence our result for  $L_1$  and  $t = 1$  from which (6.4) follows.  $\square$

**Remark 6.3.** We do not know how near the power  $m$  required in Theorem 6.2 is to the optimal for the space  $L_1$  or  $L_\infty$ , but see also Theorem 7.7.

We will now give the following crucial estimate.

**Theorem 6.4.** *For  $f \in L_p(\mathbf{R}^d)$ ,  $d \geq 2$  we have*

$$\|\Delta V_t^m f\|_p \leq \frac{\varepsilon(m)}{t^2} \|f\|_p \tag{6.5}$$

with  $\varepsilon(m) \rightarrow 0$  as  $m \rightarrow \infty$ .

**Proof.** As discussed in the proof of Theorem 6.2, it is sufficient to show that

$$\left\| \mathcal{F}^{-1} \left( |x|^2 \left( \frac{\Gamma(\frac{d}{2}) J_v(2\pi|x|)}{(\pi|x|)^v} \right)^m \right) \right\|_{L_1(\mathbf{R}^d)} \rightarrow 0 \quad \text{as } m \rightarrow \infty.$$

To show the latter, we have to show that

$$\left\| D^\mu \left\{ |x|^2 \left( \frac{\Gamma(\frac{d}{2}) J_v(2\pi|x|)}{(\pi|x|)^v} \right)^m \right\} \right\|_1 \rightarrow 0 \quad \text{as } m \rightarrow \infty$$

for the derivatives  $D^\mu$  where  $|\mu| \leq d + 1$  (including  $\mu = 0$  and  $v = \frac{d-2}{2}$ ). For  $|\mu| = \ell$ ,  $\ell \leq d + 1$  set the domain

$$D_\alpha = \left\{ x : |x| \leq \frac{1}{m^\alpha} \right\}$$

with  $\alpha$  to be chosen ( $\alpha = \frac{1}{4} \left( \frac{2d+\frac{5}{2}}{d+\frac{5}{2}} \right)$  will do).

We now write for  $\frac{d+1}{2d+3} < \alpha < \frac{1}{2}$

$$\begin{aligned} & \left\| D^\mu \left\{ |x|^2 \left( \frac{\Gamma(\frac{d}{2}) J_\nu(2\pi|x|)}{(\pi|x|)^\nu} \right)^m \right\} \right\|_{L_1(D_x)} \\ & \leq \frac{C}{m^{(d+2)\alpha}} m^{|\mu|} \left( \frac{1}{m^\alpha} \right)^{|\mu|} = o(1), \quad m \rightarrow \infty, \quad |\mu| \leq d+1, \end{aligned}$$

using the recursion relation  $\frac{d}{dt}(t^{-\nu} J_\nu(t)) = -t^{-\nu} J_{\nu+1}(t)$  and the trivial estimate  $J_{\nu+1}(t) = O(t^{\nu+1})$ ,  $t \rightarrow 0$ . For  $\frac{1}{m^\alpha} \leq |x| \leq \frac{1}{3}$  we use the estimate ( $\nu = \frac{d-2}{2}$ )

$$\begin{aligned} I &= m^{|\mu|} \left| \frac{\Gamma(\frac{d}{2})}{(\pi|x|)^\nu} J_\nu(2\pi|x|) \right|^{m-\ell} \\ &\leq m^{|\mu|} \left| \frac{\Gamma(\frac{d}{2})}{\Gamma(\frac{d-1}{2}) \Gamma(\frac{1}{2})} \int_{-1}^1 e^{i2\pi|x|s} (1-s^2)^{\frac{d-3}{2}} ds \right|^{m-\ell} \\ &\leq m^{|\mu|} \left| \frac{1}{2} (\cos 2\pi m^{-\alpha} \delta + 1) \right|^{m-\ell}, \end{aligned}$$

where  $\frac{\Gamma(\frac{d}{2})}{\Gamma(\frac{d-1}{2}) \Gamma(\frac{1}{2})} \int_{-\delta}^{\delta} (1-s^2)^{(d-3)/2} ds = \frac{1}{2}$  (recall  $\frac{\Gamma(\frac{d}{2})}{\Gamma(\frac{d-1}{2}) \Gamma(\frac{1}{2})} \int_{-1}^1 (1-s^2)^{(d-3)/2} ds = 1$ ).

Therefore,

$$I \leq m^{|\mu|} (1 - \sin^2(\pi m^{-\alpha} \delta))^{m-\ell}.$$

We choose  $\ell$  to be any fixed number  $\ell > \frac{2(d+2)}{d-1}$  from Theorem 6.2. (For example  $\ell = \left\lceil \frac{2(d+2)}{d-1} \right\rceil + 1$ .)

We can now write, as  $\sin^2 u \geq \frac{2u^2}{\pi^2}$  for  $u \leq \frac{\pi}{2}$ ,

$$I \leq m^{|\mu|} (1 - 4\delta^2 m^{-2\alpha})^{m-\ell}.$$

If  $\alpha < \frac{1}{2}$ ,  $I = o(1)$  as  $m \rightarrow \infty$ .

To complete the estimate, it is now sufficient to observe that

$$\tilde{I}(x, \nu) = \left| \frac{\Gamma(\frac{d}{2})}{(\pi|x|)^\nu} J_\nu(2\pi|x|) \right| \leq \beta < 1 \quad \text{for } |x| \geq \frac{1}{3}, \quad \nu = \frac{d-2}{2}.$$

For  $|x| \geq A$  with  $A$  big enough that estimate follows from [St-We, Lemma 3.11, p. 158]. For  $\frac{1}{3} \leq |x| \leq A$   $|\tilde{I}(x, \nu)|$  achieves a maximum at  $x_0$  as it is continuous, but for any  $x_0 \neq 0$   $\beta = |\tilde{I}(x_0, \nu)| < 1$  using the representation in [St-We, pp. 153–154].

We now choose  $\frac{d+1}{2d+3} < \alpha < \frac{1}{2}$ , and clearly  $\alpha = \frac{1}{4} \left( \frac{2d+\frac{5}{3}}{d+\frac{3}{2}} \right)$  will do. This completes the proof.  $\square$

Let us now transfer the theorems of this section to  $L_p(\mathbf{T}^d)$ .

**Theorem 6.5.** *In Theorems 6.2 and 6.4  $L_p(\mathbf{R}^d)$  can be replaced by  $L_p(\mathbf{T}^d)$ . In Theorem 6.1  $L_2(\mathbf{R}^d)$  can be replaced by  $L_2(\mathbf{T}^d)$ .*

**Proof.** As Theorems 6.2 and 6.4 are valid for  $L_\infty(\mathbf{R}^d)$ , they are valid for  $L_\infty(\mathbf{T}^d)$  since  $f \in L_\infty(\mathbf{T}^d)$  can be extended to  $f \in L_\infty(\mathbf{R}^d)$  by periodicity without changing the norm. Being valid for  $L_\infty(\mathbf{T}^d)$ , Theorems 6.2 and 6.4 follow using duality for  $L_1(\mathbf{T}^d)$ , and using the Riesz–Thorin theorem for  $L_p(\mathbf{T}^d)$ ,  $1 < p < \infty$ . For  $L_2$  the result (Theorem 6.1) follows as we have essentially the same multipliers.  $\square$

**Corollary 6.6.** *For  $f \in L_p(\mathbf{R}^d)$  or  $f \in L_p(\mathbf{T}^d)$ ,  $1 \leq p \leq \infty$  we have*

$$\|V_t f - f\|_p \approx \inf_g (\|f - g\|_p + t^2 \|\Delta g\|_p). \tag{6.6}$$

**Proof.** The proof follows that in Section 5 where we use iterations of [Di-RuI, Theorem 2.1] as seen in the Voronovskaja-type result [Di-RuI, (3.4)] in combination with Theorems 6.2 and 6.4. As the technique is almost exactly the same, we omit the details.  $\square$

**7. Smoothness of  $S_\theta f$  and  $V_t f$**

It is evident that the results on smoothness in  $L_2$ , that is, Theorem 2.1, Corollary 2.2 and Theorem 6.1 are optimal. For  $L_p$ ,  $p \neq 2$  the results we have are not optimal. However, we will show below how for  $p$  near 2 we can obtain results close to those for  $p = 2$ . It follows from Theorem 3.1 that for  $m > \frac{2(\lfloor \frac{d}{2} \rfloor + 3)}{d-2}$ ,  $\tilde{\Delta} S_\theta^m f$  exists in  $L_1$ . We will show that  $S_\theta f$  does not satisfy any smoothness property in  $L_1$ . Similar results are valid for  $V_t f$ .

We define for  $K_r(f, \tilde{\Delta}, t^{2r})_p$  given in (5.7)

$$\left\{ f : \sup_{t>0} \frac{K_r(f, \tilde{\Delta}, t^{2r})_p}{t^\alpha} < \infty \right\} \equiv L_p^\alpha(S^{d-1}), \tag{7.1}$$

and the definition does not depend on  $r$  as long as  $2r > \alpha$ .

We can now state and prove the following result.

**Theorem 7.1.** *For  $1 < p < \infty$  and  $0 < \theta \leq \frac{\pi}{2}$ ,  $f \in L_p(S^{d-1})$  implies  $S_\theta f \in L_p^{\alpha_p}(S^{d-1})$  with  $\alpha_p = (d - 2)|1 - \frac{1}{p}|$  for  $p \leq 2$ ,  $\alpha_p = (d - 2)/p$  for  $p \geq 2$  and*

$$K_r(S_\theta f, \tilde{\Delta}, t^{2r})_p \leq C \theta^{-\alpha_p} t^{\alpha_p} \|f\|_p \tag{7.2}$$

with  $C$  independent of  $t$  and  $\theta$ .



**Proof.** We set  $\eta(x) \in C^\infty[0, \infty)$  such that  $\eta(x) = \begin{cases} 1 & x \leq \frac{1}{2} \\ 0 & x \geq 1 \end{cases}$  and we denote  $\eta_j(x) \equiv \eta(\frac{x}{2^j})$ . For  $f \in L_p(S^{d-1})$ ,  $O_j f$  given by

$$O_j f \sim \sum \eta_j\left(\frac{k}{n}\right) P_k(f) \tag{7.3}$$

are multiplier operators which satisfy

$$\|O_j f\|_p \leq C \|f\|_p, \quad 1 \leq p \leq \infty, \quad j = 0, 1, \dots \tag{A}$$

and

$$E_{2^j}(f)_p \leq \|f - O_j f\|_p \leq C E_{2^{j-1}}(f)_p, \quad 1 \leq p \leq \infty, \quad j = 1, 2, \dots, \tag{B}$$

where

$$E_n(f)_p = \inf(\|f - P\|_p; \quad P \text{ spherical harmonic of degree } \leq n).$$

In fact, in our proof, any bounded sequence of multiplier operators satisfying (A) and (B) can replace  $O_j$ . (For instance, any delayed means  $V_\lambda$  generated by some Riesz means as discussed in [Ch-Di].) We now set  $T_j f \equiv O_j f - O_{j-1} f$  for  $j \geq 1$  and  $T_1 f = O_0 f$ . We show now that for  $1 \leq p \leq \infty$  and  $2r > \beta$

$$C^{-1} \sup_{j \geq 1} \frac{\|T_j f\|_p}{2^{-j\beta}} \leq \sup_{t > 0} \frac{K_r(f, \tilde{\Delta}, t^{2r})_p}{t^\beta} \leq C \sup_{j \geq 1} \frac{\|T_j f\|_p}{2^{-j\beta}}. \tag{7.4}$$

The first inequality follows as  $\|T_j f\|_p \leq A E_{2^{j-1}}(f)_p$  and  $E_n(f)_p$  is bounded by  $K_r(f, \tilde{\Delta}, n^{-2r})_p$  (see for instance [Ch-Di, (8.8)]). The second implication follows from

$$E_{2^j}(f)_p \leq \|f - O_j f\|_p \leq \sum_{\ell=j+1}^\infty \|T_\ell f\|_p \leq A 2^{-j\beta} \sup_{\ell} \left( \frac{\|T_\ell f\|_p}{2^{-\ell\beta}} \right),$$

with  $A$  depending on  $\beta$  (but not on  $j$  or  $f$ ), and the estimate of  $K_r(f, \tilde{\Delta}, t^{2r})_p$  by  $E_n(f)_p$  (see for instance [Ch-Di, (8.9)]). We now observe that for  $p = 1$

$$\|T_j(S_\theta f)\|_1 \leq \|O_j(S_\theta f)\|_1 + \|O_{j-1}(S_\theta f)\|_1 \leq 2C \|S_\theta f\|_1 \leq 2C \|f\|_1.$$

For  $p = 2$  we use  $\|T_j(S_\theta f)\|_2 \leq A E_{2^{j-1}}(S_\theta f)_2$ , and as

$$|(\theta k)^{(d-2)/2} Q_k^{(\lambda)}(\cos \theta)| \leq C \quad \text{for all } k,$$

we have  $\|(-\tilde{\Delta})^{(d-2)/4} S_\theta f\|_2 \leq 2C \theta^{-(d-2)/2} \|f\|_2$ , and hence (see [Di-II, Theorem 4.1 and also A, p. 343])  $E_{2^j}(S_\theta f)_2 \leq C_1 \theta^{-(d-2)/2} 2^{-j(d-2)/2} \|f\|_2$ .

As  $T_j$  and  $S_\theta$  are linear operators, we may use the Riesz–Thorin theorem to obtain  $\|T_j S_\theta f\|_p \leq C \theta^{-\alpha_p} 2^{-j d_p} \|f\|_p$  with  $\alpha_p = (d-2)(1 - \frac{1}{p})$  for  $1 < p < 2$ . For  $2 < q < \infty$  we get the same result by duality and  $\alpha_q = \alpha_p$  for  $\frac{1}{p} + \frac{1}{q} = 1$ .  $\square$

We cannot show that for  $1 < p < 2$  (or for  $2 < p < \infty$ ) Theorem 7.1 is best possible and it probably is not. For  $p = 2$  Theorem 7.1 is best possible and we will show

below that for  $p = 1$  (and hence by duality for  $p = \infty$ ) Theorem 7.1 cannot be improved either.

**Theorem 7.2.** *There exists  $f \in L_1(S^{d-1})$  such that  $(-\tilde{\Delta})^\alpha S_0 f \notin L_1(S^{d-1})$  for any positive  $\alpha$ .*

**Remark 7.3.** We note that using Theorem 3.1, we established that for  $m > \frac{2(\lfloor \frac{d}{2} \rfloor + 3)}{d-2}$   $\tilde{\Delta} S_0^m f \in L_1(S^{d-1})$ , and in particular if  $d \geq 10$ , we have  $\tilde{\Delta} S_0^2 f \in L_1(S^{d-1})$ , which makes Theorem 7.2 somewhat intriguing.

First we state the Nikolskii inequality for spherical polynomials.

**Lemma 7.4.** *For  $P_n$ , a spherical polynomials of degree smaller or equal to  $n$  on  $S^{d-1}$  (that is,  $P_n \in \text{span}(\bigcup_{k=0}^n H_k)$  where  $H_k = \{\psi : \tilde{\Delta}\psi = -k(k+d-2)\psi\}$ ), and  $0 < q < p \leq \infty$  we have*

$$\|P_n\|_p \leq C n^{(d-1)(\frac{1}{q}-\frac{1}{p})} \|P_n\|_q. \tag{7.5}$$

**Proof.** While we found a Ref. [Ka] for the part of the result which we need i.e. when  $q \geq 1$ , we give below a short proof of (7.5). We choose integer  $r > \frac{q}{2}$ .  $P_n(x)^r$  is a spherical polynomials of degree  $nr$ . We write

$$P_n(x)^r = \int_{S^{d-1}} P_n(y)^r \sum_{k=0}^{nr} \sum_{\ell=1}^{d_k} Y_{k,\ell}(x) Y_{k,\ell}(y) dy = \int_{S^{d-1}} P_n(y)^r G_{nr}(x \cdot y) dy,$$

where  $\{Y_{k,\ell}\}_{\ell=1}^{d_k}$  is any orthonormal system in  $H_k$ . We now have

$$\begin{aligned} \|P_n\|_\infty^r &\leq \|P_n\|_\infty^{r-q/2} \int_{S^{d-1}} |P_n(y)|^{q/2} |G_{nr}(x \cdot y)| dy \\ &\leq \|P_n\|_\infty^{r-q/2} \|P_n\|_q^{q/2} \left( \int (G_{nr}(x \cdot y))^2 dy \right)^{1/2}. \end{aligned}$$

As  $\int_{S^{d-1}} (G_{nr}(x \cdot y))^2 dy$  is independent of  $x$ , it is equal to  $\frac{1}{|S^{d-1}|} \sum_{k=0}^{nr} \dim H_k \approx n^{d-1}$ , and the proof of (7.5) when  $p = \infty$  follows.

To complete the proof we recall for  $q < p < \infty$ ,  $\|f\|_p \leq \|f\|_q^{q/p} \|f\|_\infty^{1-q/p}$ .  $\square$

**Lemma 7.5.** *If  $(-\tilde{\Delta})^\alpha f \in L_1$ , ( $\alpha < 1$ ), then  $f \in L_p$  for  $1 \leq p < p(\alpha) = \frac{d-1}{d-1-2\alpha}$ .*

**Proof.** Recall  $T_j$  from the proof of Theorem 7.1, and following [Di-II, Theorem 4.1],  $\|T_j\|_1 \leq C E_{2^{-j-1}}(f)_1 \leq C_1 2^{-2^j \alpha}$ . Using Lemma 7.4 with  $q = 1$ ,  $\|T_j\|_p \leq C_2 2^{-2^j \alpha} (2^j)^{(d-1)(1-\frac{1}{p})}$ , and if  $2\alpha > (d-1)(1-\frac{1}{p})$  or  $p < \frac{d-1}{d-1-2\alpha}$ ,

$\|\sum_{j=1}^{\infty} T_j\|_p \leq \sum \|T_j\|_p < \infty$ . As  $\|T_0\|_p \leq C\|f\|_1$  and  $\sum_{j=0}^{\infty} T_j = f$ , we complete the proof.  $\square$

**Proof of Theorem 7.2.** In view of Lemma 7.5 it is sufficient to construct a function  $f \in L_1(S^{d-1})$  such that  $S_{\theta}f \notin L_p(S^{d-1})$  for any  $p > 1$ . Our strategy is to use  $f_n(x) \geq 0$  so that  $S_{\theta}f_n \geq 0$ . We then set  $f(x) = \sum_{n=1}^{\infty} \frac{1}{n^2} f_n(x)$ , which implies  $\|S_{\theta}f\|_p \geq \frac{1}{n^2} \|S_{\theta}f_n\|_p$ , and show that  $\sup_n \frac{1}{n^2} \|S_{\theta}f_n\|_p = \infty$  for any  $p > 1$ . To do that we write

$$f_n(x) = 2^{n(d-1)} \psi_{2^{-n}}(x \cdot z_n), \quad \psi_{\delta}(t) = \begin{cases} 1 & t \geq \cos \delta, \\ 0 & t < \cos \delta. \end{cases} \tag{7.6}$$

In fact,  $z_n$  are immaterial and we can choose  $z_n = z$  if we wish. Clearly  $\|f_n\|_1 \leq C$  and  $f \in L_1(S^{d-1})$  while  $\|f_n\|_p \approx 2^{n(d-1)(1-\frac{1}{p})}$  and  $f \notin L_p(S^{d-1})$  for  $p > 1$ . We will now show that for  $\theta \geq \eta > 0$ , for example  $\theta > 10\eta > 0$  (but as  $\eta$  takes the role of  $2^{-n}$ , we could have a much larger ratio)

$$m(y : y \cdot x = \cos \theta, y \cdot z \geq \cos \eta) \geq C\eta^{d-2},$$

$$\text{if } \cos(\theta + \frac{\eta}{2}) \leq x \cdot z \leq \cos(\theta - \frac{\eta}{2}) \tag{7.7}$$

with  $C$  independent of  $\eta$ . As  $m\{x : \cos(\theta + \frac{\eta}{2}) \leq x \cdot z \leq \cos(\theta - \frac{\eta}{2})\} \approx \theta^{d-2}\eta$  the demonstration of (7.7) will complete the proof of our theorem setting  $\eta = 2^{-n}$  (and any fixed  $\theta$ ) and obtaining

$$\frac{1}{n^2} \|S_{\theta}f_n\|_p \approx \frac{1}{n^2} ((2^{-n})^{(d-2)} 2^{n(d-1)})^p \theta^{d-2} 2^{-n})^{1/p} \approx \frac{\theta^{(d-2)/p}}{n^2} 2^{n(1-\frac{1}{p})}.$$

Hence, it remains only to prove (7.7). We observe that if  $\cos(\theta - \frac{\eta}{2}) \leq x \cdot z \leq \cos(\theta + \frac{\eta}{2})$ , then there exists  $z_0$  such that  $z_0 \cdot z \geq \cos \frac{\eta}{2}$  and  $x \cdot z_0 = \cos \theta$ . This follows since in the plane containing the vectors  $x$  and  $z$ , the angle between them has to be increased or decreased by less than  $\frac{\eta}{2}$  to get  $z_0$ . The measure  $m(y : y \cdot x = \cos \theta, y \cdot z_0 \geq \cos \frac{\eta}{2}, z_0 \cdot x = \cos \theta)$  can be calculated to behave like  $\eta^{d-2}$ . Our result follows as

$$\begin{aligned} & \{y : y \cdot x = z_0 \cdot x = \cos \theta, y \cdot z_0 \geq \cos \frac{\eta}{2}\} \\ & \subset \{y : y \cdot x = \cos \theta, y \cdot z_0 \geq \cos \frac{\eta}{2}, \cos(\theta + \frac{\eta}{2}) \leq z_0 \cdot x \leq \cos(\theta - \frac{\eta}{2})\} \\ & \subset \{y : y \cdot x = \cos \theta, y \cdot z \geq \cos \eta, \cos(\theta + \eta) \leq x \cdot z \leq \cos(\theta - \eta)\}. \quad \square \end{aligned}$$

We also have the (slightly simpler to prove) analogues in  $R^d$  (or  $T^d$ ).

**Theorem 7.6.** For  $1 < p < \infty$ ,  $f \in L_p(L_p(T^d)$  or  $L_p(R^d))$  implies  $V_t f \in L_p^{\alpha_p}$  with  $\alpha_p = (d-1)|1 - \frac{1}{p}|$  if  $p \leq 2$  and  $\alpha_p = (d-1)/p$  for  $p \geq 2$  where

$$L_p^{\alpha} \equiv \left\{ f; \sup_{u>0} \frac{K_r(f, \Delta, u^{2r})_p}{u^{\alpha}} < \infty \right\}, \quad 2r > \alpha$$

and

$$K_r(f, \Delta, u^{2r})_p \equiv \inf_g (\|f - g\|_p + u^{2r} \|\Delta^r g\|_p).$$

Moreover,

$$K_r(V_t f, \Delta, u^{2r})_p \leq C t^{-\alpha_p} u^{\alpha_p} \|f\|_p. \quad (7.8)$$

**Theorem 7.7.** *There exists  $f \in L_1$  such that  $(-\Delta)^\alpha V_t f \notin L_1$  for any positive  $\alpha$ .*

**Remark 7.8.** Theorems 7.6 and 7.7 are similar and perhaps easier to prove than Theorems 7.1 and 7.2. We may prove these theorems first for  $T^d$  and deduce them for  $R^d$  or vice versa. Analogues of Lemmas 7.4 and 7.5 for  $T^d$  and trigonometric polynomials of deg  $n$  are easy to prove and perhaps better known than Lemmas 7.4 and 7.5. We did not give the proofs explicitly to avoid duplication.

**Remark 7.9.** In Theorems 7.1 and 7.2 the space  $L_p^\alpha$  appears to characterize smoothness. The same situation occurs if we use the new modulus of smoothness introduced in [Di-I], that is

$$\omega^m(f, t)_p = \sup \left\{ \left\| \Delta_\rho^m f \right\|_p; \rho x \cdot x \geq \cos t, \rho \in \text{SO}(d) \right\} \quad (7.9)$$

or if we use best spherical harmonic approximation. The above follows from the fact that for  $1 \leq p \leq \infty$

$$\omega^m(f, t)_p = O(t^\alpha) \quad \text{for } m > \alpha,$$

$$K_r(f, \tilde{\Delta}, t^{2r})_p = O(t^\alpha) \quad \text{for } 2r > \alpha$$

and

$$E_n(f)_p = O\left(\frac{1}{n^\alpha}\right)$$

are all equivalent (see [Di-III]).

It is in some more sensitive conditions that the concepts  $E_n(f)_p$ ,  $K_r(f, \tilde{\Delta}, t^{2r})_p$  and  $\omega^m(f, t)_p$  may differ.

## References

- [Bo-Cl] A. Bonami, J.L. Clerc, Sommes de Cesàro and multiplicateur de developpements en harmoniques sphériques, Trans. Amer. Math. Soc. 183 (1973) 223–262.
- [Ch-Di] W. Chen, Z. Ditzian, Best approximation and  $K$ -functionals, Acta Math. Hungar. 75 (1997) 165–208.
- [Da-Wa-Yu] F. Dai, K. Wang, C. Yu, On a conjecture of Ditzian and Runovskii, J. Approx. Theory 118 (2) (2002) 202–224.

- [Di-I] Z. Ditzian, A modulus of smoothness on the unit sphere, *J. d'Analyse Math.* 79 (1999) 189–200.
- [Di-II] Z. Ditzian, Fractional derivatives and best approximation, *Acta Math. Hungar.* 81 (1998) 323–348.
- [Di-III] Z. Ditzian, Jackson inequality on the sphere, to appear.
- [Di-Iv] Z. Ditzian, K.G. Ivanov, Strong converse inequalities, *J. d'Analyse Math.* 61 (1993) 61–111.
- [Di-RuI] Z. Ditzian, K. Runovskii, Averages and  $K$ -functionals related to the Laplacian, *J. Approx. Theory* 97 (1999) 113–139.
- [Di-RuII] Z. Ditzian, K. Runovskii, Averages on caps of  $S^{d-1}$ , *J. Math. Anal. Appl.* 248 (2000) 260–274.
- [Ka] A.I. Kamzolov, Approximation of functions on the sphere  $S^n$ , *Serdica* 10 (1984) 3–10 (in Russian).
- [Li-Ni] P.I. Lizorkin, S.M. Nikolskii, Function spaces on the sphere that are connected with approximation theory, *Mat. Zametki* 41 (1987) 509–515 (English Transl. *Math. Notes* 41 (1987) 286–291).
- [Ru] K.V. Rustamov, On equivalence of different moduli of smoothness on the sphere, *Proc. Stekelov Inst. Math.* 204 (3) (1994) 235–260.
- [St-We] E.M. Stein, G. Weiss, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton University Press, Princeton, NJ, 1971.
- [Sz] G. Szegő, *Orthogonal polynomials*, Amer. Math. Soc. Coll. Publ. 23 (1959).
- [Wa-Li] K. Wang, L. Li, *Harmonic Analysis and Approximation on the Unit Sphere*, Science Press, Beijing, China, 2000.